Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

- 1. (currently amended) An isolated nucleic acid molecule comprising:
- (a) <u>two</u> nucleotide sequences encoding a bacteriophage recombinase function;
- (b) <u>a nucleotide sequence</u> sequences encoding a bacteriophage antirecombinase function;
- (c) <u>a Ptac</u> promoter <u>sequence</u> sequences operably linked to the nucleotide sequences of (a) and (b); and
- (d) <u>a nucleotide sequence</u> sequences encoding LacI operably linked to its native promoter.
- 2. (currently amended) The nucleic acid molecule of claim 1, further comprising at least one origin of replication sequence sequences which confers confer low copy number on a vector comprising the nucleic acid molecule.
- 3. **(original)** The nucleic acid molecule of claim 2, wherein the origin of replication is temperature sensitive.
 - 4. (currently amended) An isolated nucleic acid molecule comprising:
 - (a) $\underline{\text{two}}$ nucleotide sequences encoding bacteriophage λ Red recombinase function;
 - (b) \underline{a} nucleotide sequences encoding bacteriophage λ anti-RecBCD function;
 - (c) <u>a Ptac</u> promoter <u>sequence</u> sequences operably linked to the nucleotide sequences of (a) and (b); and
 - (d) <u>a nucleotide sequence sequences</u> encoding LacI operably linked to its native promoter.
- 5. (currently amended) The nucleic acid molecule of claim 4, further comprising at least one origin of replication sequence sequences which confers eonfers low copy number on a vector comprising the nucleic acid molecule.
- 6. (original) The nucleic acid molecule of claim 5, wherein the origin of replication is temperature sensitive.

7. (currently amended) The nucleic acid molecule of any one of claims 4-6, wherein the nucleotide sequences encoding bacteriophage λ Red recombinase function comprises emprises λ exo and bet sequences.

- 8. (currently amended) The nucleic acid molecule of any one of claims 4-6, wherein the nucleotide sequences encoding λ anti-RecBCD function comprises emprises λ gam sequences.
 - 9. (currently amended) A vector comprising:
 - (a) <u>two</u> nucleotide sequences encoding a bacteriophage recombinase function;
 - (b) <u>a nucleotide sequence sequences encoding a bacteriophage anti-</u>recombinase function;
 - (c) <u>a Ptac promoter sequence</u> sequences operably linked to the nucleotide sequences of (a) and (b);
- (d) <u>a nucleotide sequence</u> sequences encoding LacI operably linked to its native promoter; and
- (e) <u>at least one origin of replication sequences which confers</u> eonfers low copy number on the vector.
- 10. (currently amended) The vector of claim 9, wherein the origin of replication sequence is sequences are temperature sensitive.
 - 11. (currently amended) A vector comprising:
 - (a) $\underline{\text{two}}$ nucleotide sequences encoding bacteriophage λ Red recombinase function;
 - (b) <u>a nucleotide sequence sequences</u> encoding bacteriophage λ anti-RecBCD function;
 - (c) <u>a Ptac promoter sequence sequences</u> operably linked to the nucleotide sequences of (a) and (b); and
 - (d) a nucleotide <u>sequence</u> sequences encoding LacI; and
 - (e) <u>at least one</u> origin of replication <u>sequence</u> sequences which <u>confers</u> eonfers low copy number on the vector.
- 12. (currently amended) The vector of claim 11, wherein the origin of replication sequence is sequences are temperature sensitive.
- 13. (currently amended) The vector of claim 12, wherein the nucleotide sequence sequences encoding-bacteriophage λ Red recombinase function comprises comprises λ exo and bet sequences.

14. (currently amended) The vector of claim 12, wherein the nucleotide sequences sequence encoding λ anti-RecBCD function comprises a comprises- λ gam sequence sequences.

- 15. (original) A recombinant organism comprising the vector of any one of claims 9-14.
 - 16. (original) The recombinant organism of claim 15, which is a bacteria.
- 17. **(original)** The recombinant organism of claim 16 which is of the genus *Escherichia*.
- 18. **(original)** The recombinant organism of claim 17, which is *Escherichia coli*.
- 19. **(original)** The recombinant organism of claim 18, which is *Escherichia coli K12*.
- 20. **(original)** The recombinant organism of claim 16 which is a pathogenic species.
- 21. **(original)** The recombinant organism of claim 20 which is a pathogenic *Escherichia coli*.
- 22. **(original)** The recombinant organism of claim 21 which is enterohemorrhagic *E. coli* (EHEC) or enteropathogenic *E. coli* (EPEC).
- 23. **(original)** The recombinant organism of claim 15 which is of the genus *Pseudomonas*.
- 24. **(original)** The recombinant organism of claim 23, which is *Pseudomonas aeruginosa*.
- 25. (original) The recombinant organism of claim 15 which is of the genus *Mycobacterium*.
- 26. (original) The recombinant organism of claim 25, which is *Mycobacterium tuberculosis*.

27. (withdrawn) A method of promoting efficient recombination of genetic material in a microorganism comprising use of the vector of any one of claims 9-14.

- 28. **(withdrawn)** The method of claim 27, wherein the genetic material is endogenous.
- 29. (withdrawn) The method of claim 27, wherein the genetic material is exogenous
- 30. (withdrawn) The method of claim 27, wherein the genetic material is derived from a prokaryote.
- 31. **(withdrawn)** The method of claim 27, wherein the genetic material is derived from a eukaryote.
- 32. **(withdrawn)** The method of claim 27, wherein the genetic material is derived from a fungi.
- 33. (withdrawn) A method for determining whether a bacterial gene is a potential drug target comprising:
 - (a) introducing a test construct into the microorganism of claim 15, wherein the test construct comprises an integrating segment flanked by recombination segments; wherein the recombination segments are homologous to the bacterial gene or surrounding sequences; and
 - (b) culturing the microorganism under conditions such that recombination between the test construct and the bacterial gene occurs; and
 - (c) assaying the microorganism for growth and/or pathogenicity or an indicator thereof,

whereby a change in growth and/or pathogenicity or an indicator thereof identifies the bacterial gene as a potential drug target.

34. (withdrawn) The method of claim 33, wherein the bacterial gene is chromosomal.

35. (withdrawn) The method of claim 33, wherein the bacterial gene is present on an endogenous plasmid.

- 36. (withdrawn) The method of claim 33, wherein the integrating segment comprises nucleotide sequences encoding a selectable marker.
- 37. (withdrawn) The method of claim 36, wherein the selectable marker is selected from the group consisting of ampicillin (Amp), kanamycin (Kan), tetracycline (Tat), and β -glycosidase (β -gal).
- 38. (withdrawn) A method of cloning a potential vaccine antigen comprising:
 - (a) introducing a substrate into the microorganism of claim 15, wherein the substrate comprises recombination segments comprising nucleotide sequences homologous to a potential vaccine antigen gene or surrounding native sequences; and
 - (b) culturing the microorganism under conditions such that recombination between the substrate and the vaccine-antigen gene sequences or surrounding native sequences occurs;

such that in vivo cloning of the vaccine antigen occurs.

- 39. (withdrawn) A vaccine comprising an antigen identified according to the method of claim 38.
- 40. **(withdrawn)** Use of the recombinant organism of claim 20 in the manufacture of a vaccine.
- 41. **(withdrawn)** A method of producing an attenuated pathogenic microorganism, comprising:
 - (a) introducing a vector of any one of claims 9-14 into a pathogenic microorganism;
 - (b) introducing a substrate into the pathogenic microorganism, wherein the substrate comprises recombination segments comprising nucleotide

sequences homologous to a gene required for pathogenicity or surrounding native sequences; and

(c) culturing the microorganism under conditions such that recombination between the substrate and the gene sequences or surrounding native sequences occurs;

such that the gene required for pathogenicity is mutated, thereby producing an attenuated pathogenic microorganism.

- 42. **(withdrawn)** An attenuated pathogenic microorganism produced according to the method of claim 41.
- 43. **(withdrawn)** A vaccine comprising an attenuated pathogenic microorganism of claim 42.